



**Hyperiforce®**  
St John's Wort tablets



## **i** Information

### Important things you need to know

- This product is used to relieve the symptoms of slightly low mood and mild anxiety.
- This product is suitable for adults over 18 years.
- **Don't take** this product if you are allergic to any of the ingredients.
- This product may affect the way some medicines work. **If you take any medicine** check the table in section 2 carefully to see if Hyperiforce is suitable for you.
- **Before you take this product:** read section 2.
- **Dosage instructions:** see section 3.
- **Talk to your doctor** if your symptoms worsen or you do not feel any benefit from this product within 6 weeks.
- Side effects are minor and rare: see section 4.

**Now read the rest of this leaflet carefully.** Keep this leaflet. You may need to read it again.

### 1. What the product is for

**Hyperiforce is a traditional herbal medicinal product used to relieve the symptoms of slightly low mood and mild anxiety. This is based on traditional use only.**

Used to help with:

- Feeling down or low
- Feeling stressed and anxious

#### Suffering from depression?

This product is for symptoms of slightly low mood and mild anxiety. If you think you are suffering from depression and have thoughts of self-harm, suicide or any other symptoms of depression you should see your doctor. Other symptoms of depression can include feeling hopeless and helpless, finding it hard to concentrate, loss of interest in your daily activities, tiredness, changes to your appetite or weight and changes in your sleeping patterns.

## **✓** **✗** Do's and Don'ts for taking this product

### 2. Before you take this product

#### Do not take this product

- If you are under 18 years of age.
- If you are allergic to any of the ingredients in this product (see section 6 Further Information).
- If you are pregnant or breast-feeding.
- If you are suffering from depression (see section 1 What the product is for).
- If you have a sensitivity to sunlight.
- If you are having light therapy (phototherapy) or any testing procedures that use light (photodiagnostic procedures).
- If you are due to have an operation in the next 10 days.
- If you are taking any of the medicines in the following table as this product may affect the way they work.
- If you are unsure if this product is suitable for you then please contact your pharmacist or our Helpline on 0845 608 5858.

|                                     |  |                                      |  |
|-------------------------------------|--|--------------------------------------|--|
| Medicines for anxiety or depression | This product is not suitable if you are taking any medicines for anxiety or depression. This includes lithium.   | Medicines to treat HIV               | Amprenavir, atazanavir, darunavir, fosamprenavir, indinavir, lopinavir, nelfinavir, ritonavir, saquinavir, tipranavir (HIV protease inhibitors) Delavirdine, efavirenz, nevirapine (HIV non-nucleoside reverse transcriptase inhibitors) |
| Hormonal contraceptives             | This product is not suitable if you are taking any hormonal contraceptives including the Pill and other forms of contraception such as hormonal implants. If you are unsure as to whether your contraception contains hormones please see your pharmacist. | Medicines for the immune system      | Ciclosporin, tacrolimus, sirolimus   |
| Emergency contraception             | The morning-after pill.  | Medicines for migraine               | Almotriptan, eletriptan, frovatriptan, naratriptan, rizatriptan, sumatriptan, zolmitriptan   |
| HRT (Hormone Replacement Therapy)   | This comes in a number of forms including tablets, patches, gels and vaginal rings.  | Barbiturates (used as tranquilisers) | Butobarbital, phenobarbital  |
| Medicines to regulate male hormones | Finasteride  | Chemotherapy                         | Dasatinib, erlotinib, etoposide, imatinib, irinotecan, mitotane, sorafenib, sunitinib Aprepitant (used to treat the side effects of chemotherapy)  |
| Medicines for heart conditions      | Amiodarone, amlodipine, digoxin, felodipine, ivabradine, nifedipine, verapamil   | Breast cancer medicine               | Exemestane   |
| Blood thinning medicines            | Acenocoumarol, warfarin  | Pain killing medicine                | Tramadol, methadone  |
| Cholesterol lowering medicines      | Atorvastatin, simvastatin  | Medicine for Parkinsons              | Rasagiline   |
| Proton pump inhibitors              | Lansoprazole, omeprazole   | Antipsychotic medicine               | Aripiprazole   |
| Medicines for an infection          | Any of the following antibiotics: clarithromycin, erythromycin, telithromycin. Any of the following antifungals: itraconazole, voriconazole  | Central Nervous System stimulants    | Methylphenidate  |
| Medicines to control diabetes       | Gliclazide   | Diuretic                             | Eplerenone   |
| Medicines for epilepsy              | This product is not suitable if you are taking any medicines for epilepsy.   | Thyroid hormones                     | Thyroxine  |
| Medicines to treat malaria          | Artemether, lumefantrine   | Medicines used to control asthma     | Theophylline   |
|                                     |  | Antihistamines                       | Fexofenadine   |





## Using

### 3. How to take this product

**Adults and the elderly:** 1 tablet three times daily.

Don't take more than the recommended dose.

Not for use in children or those under 18 years of age.

### Take special care with this product

- Avoid prolonged sunbathing or the use of sunbeds or solariums whilst taking this product. In rare cases St John's wort may cause your skin to be much more sensitive to sunlight.
- Remember to tell your doctor you are taking this product if you have to have an operation. This product should be stopped 10 days before an operation.

### Driving and using machines

- In rare cases this product may cause dizziness or tiredness. If this happens to you, do not drive or use machines.

### If you take too much

- If you take too much and feel ill, talk to your doctor. Taking too much is unlikely to be harmful.
- Your skin may be more sensitive to sunlight or UV irradiation. You should not use sunbeds or solariums and you should protect your skin from the sun for 1-2 weeks.

### If you forget to take this product

- Don't worry about the missed dose. Take the next tablet as usual.

### If you feel this product isn't working

See your doctor:

- If you do not feel an improvement within six weeks, or if you begin to feel worse.
- If your symptoms begin to interfere significantly with how you live your everyday life or if you have thoughts of self-harm or suicide.



## Side effects

### 4. Possible side effects

Like all medicines, this product can cause side effects, although not everybody gets them.

#### Minor effects

The following minor side effects can occur when using this product.

The frequency of these effects occurring is not known.

#### Digestive symptoms

- Feeling sick
- Stomach discomfort
- Digestive upsets such as diarrhoea or constipation
- Loss of appetite

These are often short-lived and should go away by themselves. However if they are troubling you, talk to your doctor or pharmacist.

#### Skin reactions

- Itching (pruritis)
- Rash
- Red, itchy spots (urticaria)
- Sunburn-like reactions after exposure to strong sunlight.

Stop taking the product if these occur.

#### Other side effects

- Headaches
- Nerve pain or tingling (neuropathy)
- Anxiety
- Mania
- Restlessness
- Dizziness
- Fatigue

If you are concerned about any side effect please talk to your doctor or pharmacist.

#### Reporting of side effects

- If you get any side effects, talk to your doctor, pharmacist or nurse. This includes any possible side effects not listed in this leaflet. You can also report side effects directly via the Yellow Card Scheme at: [www.mhra.gov.uk/yellowcard](http://www.mhra.gov.uk/yellowcard)
- By reporting side effects you can help provide more information on the safety of this medicine.



## Information

### 5. How to store this product

- Keep out of the reach and sight of children.
- Do not use this product after the expiry date on the packaging. The expiry date refers to the last day of that month.
- This product does not require any special storage conditions.
- Do not use this product if you notice a change in appearance. The tablets should be oval and brownish-green in colour.

### 6. Further Information

#### What this product contains

One tablet contains 66 mg of extract (as dry extract) from fresh St John's wort (*Hypericum perforatum* L.) flowering herb (equivalent to 170-305 mg of dried St John's wort flowering herb). Extraction solvent: Ethanol 68% V/V.

The other ingredients used for the tablet are microcrystalline cellulose, croscarmellose sodium, hydrogenated cottonseed oil and colloidal anhydrous silica.

### What this product looks like and the contents of the pack

Hyperforce tablets are oval and brownish-green in colour. Hyperforce is available in packs of 60 or 120 tablets. Not all pack sizes may be marketed.

#### Traditional Herbal Registration Holder and Manufacturer

#### Traditional Herbal Registration Holder and Batch Release:

Bioforce (UK) Ltd,  
2 Brewster Place,  
Irvine, Ayrshire, UK  
KA11 5DD

#### Manufacturer:

Bioforce AG, CH-9325,  
Roggwil Switzerland  
THR No. 13668/0014



Certification Mark

This leaflet was revised on 07/2015.

### What is St John's wort?

St. John's wort, also known as *Hypericum perforatum*, is a plant which has been used as a herbal medicine for many years. The flowers and leaves found at the top of the plant are used to make this product.

### You should also know

Bioforce runs a helpline by phone and email which can provide you with further information.

Email: [enquiries@avogel.co.uk](mailto:enquiries@avogel.co.uk)

Phone: 0845 608 5858

You can get a larger print or audio version of this leaflet.

Call this number:  
0845 608 5858.

